arXiv:0906.2638v1 [math.DG] 15 Jun 2009

Proper harmonic maps from hyperbolic Riemann surfaces into the Euclidean plane.

Antonio Alarcón[a] **and José A. Gálvez**[b]

[a]Departamento de Matemática Aplicada, Facultad de Informática, Universidad de Murcia, 30100 Espinardo, Murcia, Spain; e-mail: ant.alarcon@um.es
[b]Departamento de Geometría y Topología, Facultad de Ciencias, Universidad de Granada, 18071 Granada, Spain; e-mail: jagalvez@ugr.es

Abstract

Let Σ be a compact Riemann surface and $D_1, \ldots, D_n$ a finite number of pairwise disjoint closed disks of Σ. We prove the existence of a proper harmonic map into the Euclidean plane from a hyperbolic domain Ω containing $\Sigma \setminus \cup_{j=1}^n D_j$ and of its topological type. Here, Ω can be chosen as close as necessary to $\Sigma \setminus \cup_{j=1}^n D_j$. In particular, we obtain proper harmonic maps from the unit disk into the Euclidean plane, which disproves a conjecture posed by R. Schoen and S.T. Yau.

1 Introduction.

It is classically known that there is no proper holomorphic map from the unit disk into the complex plane. To this respect, in the more general class of harmonic maps, E. Heinz proved, in 1952, there is no harmonic diffeomorphism from the unit disk onto the Euclidean plane $\mathbb{R}^2$ [H]. However, the problem of existence of a proper harmonic map from the unit disk into the Euclidean plane remained open. Thus, in 1985, R. Schoen and S.T. Yau conjectured the non-existence of these maps [SY].

We solve that problem and, in fact, prove

Theorem 1. *Let Σ be a compact Riemann surface and $D_1, \ldots, D_n$ a family of pairwise disjoint closed disks of Σ. Consider n closed disks $D'_1, \ldots, D'_n$ such that $D'_j \subseteq intD_j$, $j = 1, \ldots, n$, then there exists a proper harmonic map*

$$\varphi : \Omega \longrightarrow \mathbb{R}^2,$$

where Ω *is a domain of the topological type of* $\Sigma\backslash\left(\cup_{j=1}^{n} D_j\right)$ *satisfying*

$$\Sigma\backslash\left(\cup_{j=1}^{n} D_j\right) \subseteq \Omega \subseteq \Sigma\backslash\left(\cup_{j=1}^{n} D'_j\right). \tag{1}$$

This result provides the existence of proper harmonic maps from a hyperbolic Riemann surface of arbitrary finite topology into the Euclidean plane. In particular, when Σ is the Riemann sphere and $n = 1$, Theorem 1 gives us the existence of proper harmonic maps from the unit disk, and hence settles negatively the conjecture above.

At this point we observe that, in general, we cannot fix the conformal structure of the domain Ω where the harmonic map is defined. However, due to the control of its topological type, this can be done when Σ is the Riemann sphere. Thus, one obtains

Corollary 1. *Let* $\Omega \subseteq \mathbb{C}$ *be any n-connected domain. Then there exist proper harmonic maps from* Ω *into the Euclidean plane.*

The paper is organized as follows. In Section 2 we establish the notation that will be used throughout this work. In addition, we prove that there is no proper harmonic map from a Riemann surface of hyperbolic type into a complete flat surface non-isometric to the Euclidean plane. This fact also motivates our study and shows that the conjecture is true when the target space is a complete flat surface non-isometric to $\mathbb{R}^2$.

We start Section 3 by proving Theorem 1, which gives the existence of proper harmonic maps from hyperbolic Riemann surfaces into the Euclidean plane. In its proof, we show a slightly more general result asserting that, under certain conditions, a harmonic map can be approximated by proper harmonic maps in any compact set of its domain. Moreover, as a consequence of the Theorem above we also prove Corollary 1.

Finally, in Section 4 we prove the following Lemma, which is the basic tool in order to prove the previous results.

Lemma 1. *Let* $\widetilde{\Sigma}$ *be a compact Riemann surface and* $K_0 \subseteq \widetilde{\Sigma}$ *a compact set. Consider two closed disks* D_1, D_2 *of* $\Sigma = \widetilde{\Sigma}\backslash int K$ *such that* $D_1 \subseteq int D_2$ *and* $K_0 \cap D_2$ *is empty. Let* $F : \Sigma\backslash int D_1 \longrightarrow \mathbb{R}^2$ *be a harmonic map and* r, R *two positive real numbers such that*

$$r < |F(p)| < R, \qquad \text{for all } p \in D_2\backslash int D_1.$$

Given three positive real numbers $\varepsilon_1, \varepsilon_2, \varepsilon_3$, *there exist a closed disk* D *and a harmonic map* $G : \Sigma\backslash int D \longrightarrow \mathbb{R}^2$ *satisfying:*

1. $D_1 \subseteq int D \subseteq D \subseteq int D_2$,

2. $|F(p) - G(p)| < \varepsilon_1$, *for all* $p \in \Sigma \backslash intD_2$,

3. $R - \varepsilon_2 < |G(p)| < R$, *for any* $p \in \partial D$,

4. $r - \varepsilon_3 < |G(p)|$, *for each* $p \in D_2 \backslash intD$.

This lemma provides, starting from a harmonic map F, a new harmonic map G defined in a slightly smaller domain. In addition, G can be chosen so that it is very close to F in a fixed compact domain $\Sigma\backslash intD_2$ and the norm of G can be made as large as needed on the boundary of its domain and is well-controlled outside $\Sigma\backslash intD_2$.

This is the key step in order to construct a suitable sequence of harmonic maps whose limit will prove Theorem 1. In the proof of Lemma 1 we shall employ some ideas used in the study of different topics such as minimal surfaces (see, for instance, [M], [AFM]) or holomorphic embeddings from Riemann surfaces into the complex 2-plane $\mathbb{C}^2$ (see [W]).

2 Preliminaries.

Let us consider the Euclidean plane $\mathbb{R}^2$ with linear coordinates (x, y) endowed with its usual induced metric $dx^2 + dy^2$.

Let Σ be a Riemann surface and $F : \Sigma \longrightarrow \mathbb{R}^2$ a map. Given an orthonormal basis $S = \{e_1, e_2\}$, we shall denote

$$F_{(1,S)} := \langle F, e_1 \rangle \quad \text{and} \quad F_{(2,S)} := \langle F, e_2 \rangle.$$

Thus, F is a harmonic map into $\mathbb{R}^2$ if and only if $F_{(j,S)} : \Sigma \longrightarrow \mathbb{R}$ are harmonic functions, $j = 1, 2$.

In our construction process we shall use that, given a harmonic map $F : \Sigma \longrightarrow \mathbb{R}^2$, a harmonic function $h : \Sigma \longrightarrow \mathbb{R}$ and an orthonormal basis $S = \{e_1, e_2\}$, then the new map $G : \Sigma \longrightarrow \mathbb{R}^2$ defined in local coordinates as

$$G_{(1,S)} = F_{(1,S)} + h \quad \text{and} \quad G_{(2,S)} = F_{(2,S)},$$

is also harmonic. This new harmonic map G can be considered as a deformation of F in the direction of e_1.

On the other hand, we shall introduce the following notation for subsets of a Riemann surface Σ:

- Given $K \subseteq \Sigma$, we denote by Σ_K the set $\Sigma\backslash int(K)$, where *int* stands for the interior of a set.

- Let D_1, D_2 be two closed disks in Σ. We shall denote $D_1 < D_2$ if $D_2 \subsetneq int(D_1)$, or equivalently, if $\Sigma_{D_1} \subsetneq int(\Sigma_{D_2})$.

We finish this Section by observing that if Σ is a Riemann surface of hyperbolic type and M is a complete flat surface which is not isometric to the Euclidean plane, then there exists no proper harmonic map from Σ into M. This fact shows a clear difference between $\mathbb{R}^2$ and the rest of complete flat surfaces.

Proposition 1. *Let Σ be a Riemann surface of hyperbolic type and M a complete flat surface non-isometric to the Euclidean plane. Then there is no proper harmonic map from Σ into M.*

Proof. Since M is a complete flat surface non-isometric to the Euclidean plane then M must be a cylinder, a Moebius band, a torus or a Klein bottle. On the other hand, assume that φ is a proper harmonic map from Σ into M. As Σ is not compact and φ is proper then M cannot be compact. That is, M must be a cylinder or a Moebius band.

Thus, if M is a complete flat cylinder then M is isometric to the Euclidean plane $\mathbb{R}^2$ under the following identification:

$$(x_1, y_1), (x_2, y_2) \text{ are identified if } x_1 = x_2 + k\,T,\ y_1 = y_2,$$

for a fixed real number $T > 0$, and $k \in \mathbb{Z}$.

Analogously, if M is a complete flat Moebius band then M is isometric to the Euclidean plane $\mathbb{R}^2$ under the identification:

$$(x_1, y_1), (x_2, y_2) \text{ are related if } x_1 = x_2 + k\,T,\ y_1 = (-1)^k y_2,$$

for a fixed real number $T > 0$, and $k \in \mathbb{Z}$.

In both cases, let us consider the function $h : M \longrightarrow \mathbb{R}$ given by the distance from a point to the compact geodesic $\mathbb{R} \times \{0\}$ on M, that is, $h(x, y) = |y|$. Then $\Sigma_0 = (h \circ \varphi)^{-1}(0)$ is compact and $h \circ \varphi : \Sigma \longrightarrow \mathbb{R}$ is a proper function which is smooth and harmonic on $\Sigma \backslash \Sigma_0$. Thus, using [G, Corollary 7.7], Σ would be parabolic.

Therefore, there is no proper harmonic map from a hyperbolic Riemann surface into M. □

3 Existence of proper harmonic maps

This Section is devoted to the construction of proper harmonic maps from hyperbolic domains of a compact Riemann surface Σ into the Euclidean plane and the study of their properties.

Thus, we start by proving our main result about the existence of proper harmonic maps from hyperbolic Riemann surfaces into $\mathbb{R}^2$. This proof fundamentally relies on Lemma 1 which will be showed in Section 4.

Proof of Theorem 1. Throughout this proof we show a more general result which will be used for our density result in Corollary 2. That is, we prove that given any harmonic map

$$F : \Sigma \backslash \left(\cup_{j=1}^{n} int(D'_j) \right) \longrightarrow \mathbb{R}^2$$

and a real number $\delta_0 > 0$, there exist an open domain Ω and a proper harmonic map $\varphi : \Omega \longrightarrow \mathbb{R}^2$ such that

$$|F(p) - \varphi(p)| < \delta_0, \qquad \text{for all } p \in \Sigma \backslash \left(\cup_{j=1}^{n} int(D_j) \right), \tag{2}$$

where Ω is of the topological type of $\Sigma \backslash \left(\cup_{j=1}^{n} D_j \right)$ and satisfies (1).

For that, we shall prove that fixed j_0, there exist a domain Ω_{j_0} and a proper harmonic map $\varphi_{j_0} : \Omega_{j_0} \longrightarrow \mathbb{R}^2$ such that

$$\left| \frac{1}{n}\, F(p) - \varphi_{j_0}(p) \right| < \delta := \frac{\delta_0}{n}, \qquad \text{for all } p \in \Sigma \backslash \left(\cup_{j=1}^{n} int(D_j) \right), \tag{3}$$

where

$$\Sigma \backslash \left(int(D_{j_0}) \cup (\cup_{j \neq j_0} int(D'_j)) \right) \subseteq \Omega_{j_0} \subseteq \Sigma \backslash \left(\cup_{j=1}^{n} int(D'_j) \right)$$

and $\Omega_{j_0} \cap int(D_{j_0})$ is an open annulus.

In such a case, the new harmonic map $\varphi = \varphi_1 + \ldots + \varphi_n$ is well defined in the open domain $\Omega = \cap_{j=1}^{n} \Omega_j$ and proper. Therefore, φ proves the Theorem and also inequality (2).

Up to a translation, we can assume the origin is not contained in the compact set $F(\cup_{j=1}^{n} D_j \backslash \cup_{j=1}^{n} int(D'_j))$. Thus, once j_0 is fixed, there exist two positive real numbers r_0, R_0 such that

$$r_0 < \frac{1}{n}\, |F(p)| < R_0, \qquad \text{for all } p \in \Sigma_{D'_{j_0}} \cap D_{j_0}.$$

Let us consider a decreasing sequence of positive real numbers η_k such that $\sum_{k \geq 1} \eta_k \leq 1/2$ and denote $R_k = R_0 + k$. Let us also call $D_1^0 = D'_{j_0}$, $D_2^0 = D_{j_0}$, $F_0 = (1/n) F$ and $\Sigma' = \Sigma \backslash int(\cup_{j \neq j_0} D'_j)$. Our first goal is to prove that there exist two sequences of closed disks D_1^k, D_2^k and a sequence of harmonic maps $F_k : \Sigma_{D_1^k} \longrightarrow \mathbb{R}^2$ satisfying

(p1) $D_2^k < D_2^{k+1} < D_1^{k+1} < D_1^k$.

(p2) $|F_{k+1}(p) - F_k(p)| < \delta\, \eta_{k+1}$, for all $p \in \Sigma'_{D_2^k}$.

(p3) $R_{k+1} - \eta_{k+1} < |F_{k+1}(p)| < R_{k+1}$, for any $p \in \Sigma'_{D_1^{k+1}} \cap D_2^{k+1}$.

(p4) $R_k - \eta_k - \eta_{k+1} < |F_{k+1}(p)|$, for each $p \in \Sigma'_{D_1^{k+1}} \cap D_2^k$.

Using Lemma 1 for the Riemann surface Σ, the compact domain $\cup_{j \neq j_0} D'_j$ and the harmonic map F_0, we have the existence of a closed disk D_1^1 such that

- $D_2^0 < D_1^1 < D_1^0$.
- $|F_1(p) - F_0(p)| < \delta\, \eta_1$, for all $p \in \Sigma'_{D_2^0}$.
- $(R_0 + 1) - \eta_1 < |F_1(p)| < R_0 + 1$, for any $p \in \partial D_1^1$.

Thus, from the last inequality, one gets the existence of a closed disk D_2^1 such that $D_2^0 < D_2^1 < D_1^1$ and

$$(R+1) - \eta_1 < |F_1(p)| < R + 1, \qquad \text{for any } p \in \Sigma'_{D_1^1} \cap D_2^1.$$

Therefore, the properties (p1),(p2) and (p3) are satisfied. Observe that the property (p4) has no sense for this first step.

Now, let us assume the sequences D_1^k, D_2^k, F_k are well defined until $k = m$ and satisfy properties (p1)–(p4). Then we define the following elements of the sequences for $k = m+1$ as follows. Note that property (p4) will not be used in the construction of the remaining terms of the sequence.

Using again Lemma 1 and bearing in mind that (p3) is satisfied for $k = m$, we obtain the existence of a closed disk D_1^{m+1} such that

- $D_2^m < D_1^{m+1} < D_1^m$.
- $|F_{m+1}(p) - F_m(p)| < \delta\, \eta_{m+1}$, for all $p \in \Sigma'_{D_2^m}$.
- $R_{m+1} - \eta_{m+1} < |F_{m+1}(p)| < R_{m+1}$, for any $p \in \partial D_1^{m+1}$.
- $(R_m - \eta_m) - \eta_{m+1} < |F_{m+1}(p)|$, for each $p \in \Sigma'_{D_1^{m+1}} \cap D_2^m$.

So, since $D_2^m < D_1^{m+1}$ and from the inequality satisfied for the points in ∂D_1^{m+1}, one obtains the existence of a closed disk D_2^{m+1} such that $D_2^m < D_2^{m+1} < D_1^{m+1}$ and

$$R_{m+1} - \eta_{m+1} < |F_{m+1}(p)| < R_{m+1}, \qquad \text{for any } p \in \Sigma'_{D_1^{m+1}} \cap D_2^{m+1}.$$

Hence, the properties (p1)–(p4) are also satisfied for $k = m+1$.

Let us now define the desired harmonic map φ_{j_0}. For that, we first define $\Omega_{j_0} = \cup_{k\geq 0}\Sigma'\backslash D_2^k$. Observe that $\Sigma'\backslash D_2^k \subseteq \Sigma'\backslash D_2^{k+1}$ and $int(D_2^k)\backslash D_2^{k+1}$ is an open annulus. So, an elementary topological argument gives us $\Omega_{j_0} \cap int(D_2^0)$ is an open annulus.

In addition, let us see that for any compact set $K \subseteq \Omega_{j_0}$ we have that $F_{k|K}$ is a Cauchy sequence of harmonic maps. In such a case, the limit map φ_{j_0} is harmonic and well defined in Ω_{j_0}.

Thus, let us fix the compact set $K \subseteq \Omega_{j_0}$. Then, there exists a non-negative entire number k_0 such that $K \subseteq \Sigma'_{D_2^{k_0}}$. Hence, using (p2), for $k_1 > k_2 \geq k_0$

$$|F_{k_1}(p) - F_{k_2}(p)| < \delta \sum_{j=k_2+1}^{k_1} \eta_j, \qquad \text{for all } p \in K, \tag{4}$$

and, so, $F_{k|K}$ is a Cauchy sequence since $\sum_{j\geq 1}\eta_j \leq 1/2 < \infty$.

Taking limits in (4), we also obtain that (3) is satisfied, since if $p \in \Sigma'_{D_2^0}$ then

$$|F_0(p) - \varphi_{j_0}(p)| \leq \delta \sum_{k\geq 1} \eta_k < \delta.$$

Finally, we prove that φ_{j_0} is proper. Let m be a positive entire and consider a point $p \in \Sigma'_{D_2^{m+1}} \cap D_2^m$, then from (4) and (p4) one has

$$|\varphi_{j_0}(p) - F_{m+1}(p)| \leq \delta \sum_{k\geq m+2} \eta_k < \delta,$$
$$|F_{m+1}(p)| > R_m - \eta_m - \eta_{m+1} > R_m - 1 = R_0 + m - 1.$$

And so $|\varphi_{j_0}(p)| > m + (R_0 - \delta - 1)$.

In particular, if $q \in \Omega_{j_0} \cap D_2^m$ then there exists $m' \geq m$ such that $q \in \Sigma'_{D_2^{m'+1}} \cap D_2^{m'}$. Thereby,

$$|\varphi_{j_0}(q)| > m + (R_0 - \delta - 1), \qquad \text{for all } q \in \Omega_{j_0} \cap D_2^m.$$

Hence, for any real number $N > 0$ there exists a positive entire m such that the set $\{q \in \Omega_{j_0} : \ |\varphi_{j_0}(q)| \leq N\}$ is contained in the compact set $\Sigma'_{D_2^m}$. Therefore, φ_{j_0} is proper as we wanted to show. □

When we consider Σ as the Riemann sphere in Theorem 1, and remove only one closed disk, we get proper harmonic maps from a hyperbolic simply connected domain Ω into $\mathbb{R}^2$. Thus, Ω must be conformal to the unit disk, obtaining proper harmonic maps from the unit disk into the Euclidean plane.

Remark 1. *In general, the proper harmonic maps given by Theorem 1 could be non-surjective. However, though we shall not worry about this problem, following the Proof of Lemma 1, it is always possible to find a proper harmonic map φ given by Theorem 1 which is onto.*

It is important to note that in Theorem 1 we control the topological type of the domain Ω where the harmonic map is defined. However, we cannot fix the conformal type in general. But, this can be done when we start with a domain in the Riemann sphere.

So, as an immediate consequence of the previous Theorem, we prove that if Ω is an n-connected domain in $\mathbb{C}$, then there exist proper harmonic maps from Ω into $\mathbb{R}^2$.

Proof of Corollary 1. Let Ω be an n-connected domain in $\mathbb{C}$.

If $\mathbb{C}\backslash\Omega$ is a finite number of points, the result is obvious. Otherwise, Ω is conformal to a bounded domain whose boundary is made by a finite number of pairwise disjoint circles $C_1, \ldots, C_k$ and a finite number of points $p_{k+1}, \ldots, p_n$.

Let D_j be the closed disk in the Riemann sphere $\overline{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$, such that D_j does not intersect Ω and whose boundary is C_j. Since, $\overline{\mathbb{C}}\backslash D_j$ is conformal to the unit disk, using Theorem 1, there exists a proper harmonic map $\varphi_j : \overline{\mathbb{C}}\backslash D_j \longrightarrow \mathbb{R}^2$.

In addition, for the points p_j we take φ_j as a holomorphic function from $\overline{\mathbb{C}}\backslash\{p_j\}$ with a pole at p_j.

Thus, it is easy to see that the new harmonic map $\varphi = \sum_{j=1}^{n} \varphi_j : \Omega \longrightarrow \mathbb{R}^2$ is proper. □

On the other hand, observe from (2) that every harmonic map defined on a Riemann surface minus n open disks into $\mathbb{R}^2$ can be approximated by proper harmonic maps in every compact set contained in the interior of the domain. That is, we have really showed the following density result.

Corollary 2. *Let Σ be a Riemann surface and $D_1, \ldots, D_n$ a family of pairwise disjoint closed topological disks of Σ. Consider a harmonic map $F : \Sigma\backslash\cup_{j=1}^{n} int(D_j) \longrightarrow \mathbb{R}^2$. Then for any compact set $K \subseteq \Sigma\backslash \cup_{j=1}^{n} D_j$ and $\delta_0 > 0$, there exist a hyperbolic domain $\Omega \subseteq \Sigma$ containing K and a proper harmonic map $\varphi : \Omega \longrightarrow \mathbb{R}^2$ such that*

$$|\varphi(p) - F(p)| < \delta_0, \qquad \text{for all } p \in K.$$

Notice Schoen and Yau related their conjecture with the problem of non-existence of a hyperbolic minimal surface in $\mathbb{R}^3$ which properly projects into a plane [SY]. However, since there exist proper harmonic maps from hyperbolic Riemann surfaces, the problem of finding such a hyperbolic minimal surface remains open.

Moreover, Schoen and Yau proved that there is no harmonic diffeomorphism from the unit disk onto a complete surface with non-negative Gauss curvature [SY], extending the result of Heinz for the Euclidean plane [H]. To this respect, it would be interesting to study the existence of proper harmonic maps from hyperbolic Riemann surfaces into a complete surface with non-negative Gauss curvature.

Finally, it should be also mentioned that other Picard type problems for harmonic diffeomorphisms from the complex plane on Hadamard surfaces have been recently studied [CR], [GR], though the used techniques in that context are completely different.

4 Proof of Lemma 1.

We fix a real number $\varepsilon_0 > 0$ which satisfies some inequalities in terms of the initial data $r, R, \varepsilon_1, \varepsilon_3$ and whose dependence will be made clear along this proof.

We take a closed disk D_3 on Σ such that $D_2 < D_3 < D_1$ and a set of points $\{p_1, \ldots, p_n\}$ in the open set $int(\Sigma_{D_3}) \backslash \Sigma_{D_2}$ satisfying:

(a1) There exists a closed disk D_4 such that $\{p_1, \ldots, p_n\} \subseteq \partial D_4$ with $D_2 < D_4 < D_3$. Moreover, we can assume these points are ordered along the circle ∂D_4.

(a2) There exist open disks $B_j \subseteq int(\Sigma_{D_3}) \backslash \Sigma_{D_2}$ such that $p_j, p_{j+1} \in B_j$ and

$$|F(p) - F(q)| < \varepsilon_0, \qquad \text{for each } p, q \in B_j.$$

Here and from now on, we shall suppose $p_{n+1} = p_1$.

(a3) The well oriented orthonormal basis $S_j = \{\frac{F(p_j)}{|F(p_j)|}, i\,\frac{F(p_j)}{|F(p_j)|}\}$ satisfy

$$\left|\frac{F(p_j)}{|F(p_j)|} - \frac{F(p_{j+1})}{|F(p_{j+1})|}\right| < \frac{\varepsilon_0}{3(R-r)}, \qquad j = 1, \ldots, n.$$

Observe that these points can be chosen by continuity of F in the compact set Σ_{D_1}.

Now, let us take a Riemannian metric $\langle , \rangle$ on Σ which is compatible with the conformal structure and consider a real number $\delta > 0$ such that:

- $D_5 = D_4 \cup \left(\cup_{j=1}^{n} D(p_j, \delta)\right)$ is a topological closed disk. Here, $D(p, \delta)$ denotes the closed geodesic disk center at p and radius δ for the metric $\langle , \rangle$.
- $D(p_j, \delta) \cup D(p_{j+1}, \delta) \subseteq B_j$, for all $j = 1, \ldots, n$.
- $D(p_j, \delta) \cap D(p_k, \delta)$ is empty for any $j \neq k$.
- $F_{(1,S_j)}(p) > r$ for each $p \in D(p_j, \delta)$.

Let $\widehat{\zeta}_j : \Sigma_{D_1}\backslash\{p_j\} \longrightarrow \mathbb{C}$ be a holomorphic function with a simple pole at p_j. Its existence is clear if Σ is a topological sphere and assured by the Noether gap Theorem when the genus of Σ is positive.

Up to multiplication of $\widehat{\zeta}_j$ by a complex number, if necessary, there exist a point $q_j \in \partial D(p_j,\delta) \cap int(\Sigma_{D_4})$ and a simple curve $\beta_j : [0,1] \longrightarrow D(p_j,\delta)$ such that

- $\beta_j(0) = q_j$ and $\beta_j(1) = p_j$.
- $\beta_j(]0,1]) \subseteq int D(p_j,\delta)$.
- $\widehat{\zeta}_j(\beta_j(t))$ is a positive real number for all $t \in [0,1[$.
- $\lim_{t\to 1} \widehat{\zeta}_j(\beta_j(t)) = \infty$.

Let us denote by K_j to the compact set $\Sigma_{D_1}\backslash int(D(p_j,\delta))$ and by ζ_j to the real part of $\widehat{\zeta}_j$. Then, we can chose $\kappa > 0$ small enough in order to satisfy

$$\kappa \max\{|\zeta_j(p)| : \ p \in K_j\} < \min\left\{\frac{\varepsilon_0}{n}, 3\,(R-r)\right\}, \qquad j = 1,\ldots,n. \tag{5}$$

And we call $a_j = \beta_j(t_j)$ where

$$t_j = \min\{t \in]0,1[: \ \kappa\, \widehat{\zeta}_j(\beta_j(t)) = \kappa\, \zeta_j(\beta_j(t)) = 3(R-r)\}.$$

Now, we construct a family of harmonic maps $h_j : U_j \longrightarrow \mathbb{R}^2$, where $U_j = \Sigma_{D_1}\backslash\{p_1,\ldots,p_j\}$. These harmonic maps are defined as follows

- $h_0 = F$.
- In the basis S_j we have

$$h_{j\,(1,S_j)} = h_{j-1\,(1,S_j)} + \kappa\,\zeta_j, \qquad h_{j\,(2,S_j)} = h_{j-1\,(2,S_j)}.$$

Lemma 2. *These new harmonic maps satisfy the following properties*

(b1) $|h_j(p) - h_{j-1}(p)| < \varepsilon_0/n$, *for any* $p \in U_j\backslash int(D(p_j,\delta))$.

(b2) $|h_j(a_j) - h_{j-1}(a_{j-1})| < 4\,\varepsilon_0$.

(b3) $|h_n(a_j) - h_n(a_{j+1})| < 6\,\varepsilon_0$.

(b4) $|h_n(a_j)| > 2R - r$.

Proof. In order to obtain the inequality (b1), we have from (5)

$$|h_j(p) - h_{j-1}(p)| = |\kappa\,\zeta_j(p)| < \varepsilon_0/n.$$

Let us write $S_j = \{e_1^j, e_2^j\}$, that is, $e_1^j = F(p_j)/|F(p_j)|$ and $e_2^j = i\,F(p_j)/|F(p_j)|$, then one has for $j \geq 2$

$$\begin{aligned} |h_j(a_j) - h_{j-1}(a_{j-1})| &= |h_{j-1}(a_j) + \kappa\,\zeta_j(a_j)e_1^j - h_{j-2}(a_{j-1}) - \kappa\,\zeta_{j-1}(a_{j-1})e_1^{j-1}| \\ &\leq |h_{j-1}(a_j) - F(a_j)| + |F(a_j) - F(a_{j-1})| \\ &\quad + |F(a_{j-1}) - h_{j-2}(a_{j-1})| + 3(R-r)|e_1^j - e_1^{j-1}|. \end{aligned}$$

In addition, from (b1), $|h_{j-1}(a_j) - F(a_j)|$ and $|h_{j-2}(a_{j-1}) - F(a_{j-1})|$ are less than ε_0. On the other hand, using (a2), $|F(a_j) - F(a_{j-1})| < \varepsilon_0$ since $a_{j-1}, a_j \in B_{j-1}$. Hence, from (a3), the inequality (b2) is proved.

In order to obtain (b3) we observe

$$\begin{aligned} |h_n(a_j) - h_n(a_{j+1})| &\leq |h_n(a_j) - h_j(a_j)| + |h_j(a_j) - h_{j+1}(a_{j+1})| \\ &\quad + |h_{j+1}(a_{j+1}) - h_n(a_{j+1})| \; < \; \varepsilon_0 + 4\varepsilon_0 + \varepsilon_0 \; = \; 6\varepsilon_0, \end{aligned}$$

where we have used (b1) and (b2).

Moreover, from (b1),

$$\begin{aligned} |h_n(a_j)| &> |h_j(a_j)| - \varepsilon_0 \\ &= \sqrt{\Big(h_{j-1_{(1,S_j)}}(a_j) + \kappa\,\zeta_j(a_j)\Big)^2 + h_{j-1_{(2,S_j)}}(a_j)^2} - \varepsilon_0 \\ &\geq |h_{j-1_{(1,S_j)}}(a_j) + 3(R-r)| - \varepsilon_0, \end{aligned}$$

since $\kappa\,\zeta_j(a_j) = 3(R-r)$.

We also have from (a2) and (b1)

$$\begin{aligned} |h_{j-1_{(1,S_j)}}(a_j) - F_{(1,S_j)}(p_j)| &\leq |h_{j-1}(a_j) - F(p_j)| \\ &\leq |h_{j-1}(a_j) - F(a_j)| + |F(a_j) - F(p_j)| \; < \; 2\varepsilon_0. \end{aligned}$$

As $F_{(1,S_j)}(p_j) = |F(p_j)| > r$ by hypothesis, then $h_{j-1_{(1,S_j)}}(a_j) > r - 2\varepsilon_0$ and so

$$|h_n(a_j)| > 3R - 2r - 3\varepsilon_0 > 2R - r.$$

Here, we use that ε_0 can be initially chosen with $R - r - 3\varepsilon_0 > 0$. □

Let $\widetilde{C}_j$ be a simple regular curve contained in $int(D(p_j, \delta))$ which is transversal to $\beta_j([0,1])$ at a_j. Consider a small closed connected neighborhood C_j of the curve $\widetilde{C}_j$ at the point a_j, such that each connected component of $C_j \backslash \{a_j\}$ lies in one side of $\beta_j([0,1])$. Moreover, C_j is chosen small enough in order to satisfy (b5) $|h_j(p) - h_j(a_j)| < \varepsilon_0$ for all $p \in C_j$.

Consider a topological closed strip G_j contained in $D(p_j, \delta)$ such that $\beta_j(]0, t_j[) \subseteq int(G_j)$, and bounded by C_j, $\partial D(p_j, \delta)$ and two closed simple curves lying at each side of $\beta_j([0, t_j])$ and joining a point of $\partial D(p_j, \delta)$ with a point of C_j (see Figure 1).

In addition, we take G_j small enough such that

(b6) $|h_j(p) - h_{j-1}(p) - k\,\zeta_j(p) e_1^j| < \varepsilon_0$ for all $p \in G_j$.

(b7) $0 < k\,\zeta_j(p) < 3(R - r) + \varepsilon_0$ for each $p \in G_j$.

Note that G_j can be chosen in order to satisfy these two properties. For (b6), it is sufficient to observe that $h_j(p) = h_{j-1}(p) + k\,\zeta_j(p) e_1^j$ for any point p on $\beta_j([0, 1[)$. And (b7) is a consequence of the definition of a_j and ζ_j.

Now, we consider the compact set

$$K = \big(\Sigma_{D_4} \backslash \cup_{j=1}^n int(D(p_j, \delta))\big) \cup \big(\cup_{j=1}^n G_j\big)$$

and D_6 the closed disk $\Sigma \backslash int K$. That is, $K = \Sigma_{D_6}$.

For this new set we observe that

- $\beta_j([0, t_j]) \subseteq \Sigma_{D_6}$,
- $D_2 < D_6$,
- the harmonic maps h_j are well defined in Σ_{D_6},
- $\Sigma_{D_6} \cap int(D(p_j, \delta)) \subseteq G_j \subseteq \Sigma_{D_6} \cap D(p_j, \delta)$.

Finally, we take a closed disk D_7 such that $D_6 < D_7$ and the harmonic maps h_j are well defined in Σ_{D_7}.

Until now, we have deformed the initial harmonic map F obtaining a new harmonic map h_n. These harmonic maps are close in the compact domain Σ_{D_2} (property (b1)). And the norm of h_n is bigger than $2R - r > R$ for the family of points $a_1, \ldots, a_n$ (property (b4)). Thus, we have obtained a harmonic map which satisfies part of the conditions of Lemma 1, but only near the points a_j.

Our goal is to deform h_n in order to obtain the harmonic map which proves the Lemma. For that, we shall construct, from h_n, a sequence of n harmonic maps. Each harmonic map will be close to the previous one in a large compact domain and only modified along a neighborhood of a curve joining a_{j-1} and a_j. The last harmonic map of this sequence will be the one which solves Lemma 1.

Let us denote by Q_j the connected component of $\overline{\partial D_6 \backslash (C_j \cup C_{j+1})}$ which does not intersect C_k for all $k \neq j, j+1$. These closed curves satisfy (see Figure 1)



Figure 1.

- $Q_j \cap Q_k$ is empty for each $j \neq k$,
- $Q_j \subseteq B_j$,
- $Q_j \cap D(p_k, \delta)$ is empty for each $k \neq j, j+1$,
- $\partial D_6 = \left(\cup_{j=1}^{n} Q_j\right) \cup \left(\cup_{j=1}^{n} C_j\right)$.

Consider an open neighborhood $\widehat{C}_j$ of C_j such that

$$|h_n(p) - h_n(a_j)| < 3\,\varepsilon_0, \qquad \text{for all } p \in \widehat{C}_j \cap \Sigma_{D_5}. \tag{6}$$

This can be done by continuity, because for any $p \in C_j$

$$|h_n(p) - h_n(a_j)| \leq |h_n(p) - h_j(p)| + |h_j(p) - h_j(a_j)| + |h_j(a_j) - h_n(a_j)| < 3\,\varepsilon_0,$$

where we have used (b1) and (b5).

We also consider the neighborhoods of Q_j

$$Q_j(L) = \{p \in \Sigma_{D_1} : \ \operatorname{dist}(p, Q_j) \leq L\},$$

and fix $L_0 > 0$ small enough in order to satisfy similar properties to Q_j, that is,

- $Q_j(L_0) \cap Q_k(L_0)$ is empty for any $j \neq k$,
- $Q_j(L_0) \subseteq B_j \cap int(\Sigma_{D_7})$,

- $Q_j(L_0) \cap D(p_k, \delta)$ is empty for each $k \neq j, j+1$.

We are now ready to define a sequence of harmonic maps $g_j : \Sigma \backslash D_7 \longrightarrow \mathbb{R}^2$, $j = 0, \ldots, n$. For that, we choose some basis which will indicate the direction of the deformation. So, let $\overline{S}_j = \{\overline{e}^1_j, \overline{e}^2_j\}$, with

$$\overline{e}^2_j = \frac{h_n(a_j)}{|h_n(a_j)|}, \quad \overline{e}^1_j = -i\, \overline{e}^2_j.$$

Let $g_0 = h_n$ and let us define g_j from g_{j-1} as follows. Let

$$\tau_j = 2R - r - \min \left\{ g_{j-1_{(1,\overline{S}_j)}}(p) : \ p \in Q_j \right\}.$$

By Royden's Theorem [R, Theorem 10], there exists a holomorphic function $\widehat{\xi}_j : \Sigma_{D_7} \longrightarrow \mathbb{C}$ such that

- $|\widehat{\xi}_j(p)| < \varepsilon_0/n$, for any $p \in \Sigma_{D_6} \backslash int(Q_j(L_0))$,
- $|\widehat{\xi}_j(p) - \tau_j| < \varepsilon_0/n$, for all $p \in Q_j(L_0/2)$.

If we take the harmonic function $\xi_j : \Sigma_{D_7} \longrightarrow \mathbb{R}$ given by the real part of $\widehat{\xi}_j$, then

- $|\xi_j(p)| < \varepsilon_0/n$, for any $p \in \Sigma_{D_6} \backslash int(Q_j(L_0))$,
- $|\xi_j(p) - \tau_j| < \varepsilon_0/n$, for all $p \in Q_j(L_0/2)$.

Thus, we define

$$g_{j_{(1,\overline{S}_j)}} = g_{j-1_{(1,\overline{S}_j)}} + \xi_j, \qquad g_{j_{(2,\overline{S}_j)}} = g_{j-1_{(2,\overline{S}_j)}}.$$

The harmonic maps g_j satisfy the following properties

(c1) $|g_j(p) - g_{j-1}(p)| < \varepsilon_0/n$, for any $p \in \Sigma_{D_6} \backslash int(Q_j(L_0))$.

(c2) $|g_j(p) - g_{j-1}(p) - \tau_j\, \overline{e}^j_1| = |\xi_j(p) - \tau_j| < \varepsilon_0/n$, for all $p \in Q_j(L_0/2)$.

Finally, we define the harmonic map $G : \Sigma_{D_6} \longrightarrow \mathbb{R}^2$ as $G = g_n$. Therefore, Lemma 1 will be a consequence of the next result by shrinking the domain Σ_{D_6}.

Lemma 3. *The harmonic map G satisfies*

1. $|F(p) - G(p)| < \varepsilon_1$, *for all* $p \in \Sigma_{D_2}$,
2. $|G(p)| > R$, *for any* $p \in \partial D_6$,

3. $|G(p)| > r - \varepsilon_3$*, for each* $p \in \Sigma_{D_6} \cap D_2$*.*

Proof. The first inequality is a consequence of (b1) and (c1), since if $p \in \Sigma_{D_2}$

$$|F(p) - G(p)| \le |g_n(p) - g_0(p)| + |h_n(p) - h_0(p)| < 2\,\varepsilon_0,$$

where we can choose initially $\varepsilon_0 < \varepsilon_1/2$.

In order to prove the second inequality, we distinguish four cases:

(i) Let $p \in \widehat{C}_j \cap Q_j(L_0)$. Using (c1), (6) and (b4), one gets

$$\begin{aligned} |G(p)| &= |g_n(p)| \;>\; |g_j(p)| - \varepsilon_0 \;\ge\; |g_{j(2,\overline{S}_j)}(p)| - \varepsilon_0 \;=\; |g_{j-1(2,\overline{S}_j)}(p)| - \varepsilon_0 \\ &\ge |g_{0(2,\overline{S}_j)}(p)| - 2\varepsilon_0 \;=\; |h_{n(2,\overline{S}_j)}(p)| - 2\varepsilon_0 \;>\; |h_{n(2,\overline{S}_j)}(a_j)| - 5\varepsilon_0 \\ &= |h_n(a_j)| - 5\varepsilon_0 \;>\; 2R - r - 5\varepsilon_0 > R. \end{aligned}$$

Where ε_0 initially satisfies $R - r - 5\varepsilon_0 > 0$.

(ii) Let $p \in \widehat{C}_j \cap Q_{j-1}(L_0)$. Reasoning as in the previous case and using (b3)

$$\begin{aligned} |G(p)| &> |h_{n(2,\overline{S}_{j-1})}(a_j)| - 5\varepsilon_0 \;>\; |h_{n(2,\overline{S}_{j-1})}(a_{j-1})| - 11\varepsilon_0 \;=\; |h_n(a_{j-1})| - 11\varepsilon_0 \\ &> 2R - r - 11\varepsilon_0 \;>\; R, \end{aligned}$$

with $R - r - 11\varepsilon_0 > 0$ a priori.

(iii) Let $p \in \widehat{C}_j \backslash (\cup_{k=1}^n Q_k(L_0))$. From (c1) and (6), one obtains

$$|G(p)| \;>\; |g_0(p)| - \varepsilon_0 \;=\; |h_n(p)| - \varepsilon_0 \;>\; |h_n(a_j)| - 4\varepsilon_0 \;>\; 2R - r - 4\varepsilon_0 \;>\; R.$$

(iv) Let $p \in Q_j \backslash \left(\cup_{k=1}^n \widehat{C}_k\right)$. Then, from (c1) and (c2)

$$\begin{aligned} |G(p)| &> |g_j(p)| - \varepsilon_0 \;\ge\; |g_{j(1,\overline{S}_j)}(p)| - \varepsilon_0 \;>\; |g_{j-1(1,\overline{S}_j)}(p) + \tau_j| - 2\varepsilon_0 \\ &\ge g_{j-1(1,\overline{S}_j)}(p) + \tau_j - 2\varepsilon_0 \;\ge\; 2R - r - 2\varepsilon_0 > R. \end{aligned}$$

Finally, we prove the third inequality, that is, $|G(p)| > r - \varepsilon_3$. For that, we need to distinguish five different cases, depending on the region of Σ_{D_6} where p is located.

(i) Let $p \notin \left(\cup_{j=1}^n D(p_j, \delta)\right) \cup \left(\cup_{j=1}^n Q_j(L_0)\right)$. Using (c1) and (b1)

$$|G(p)| \;>\; |h_n(p)| - \varepsilon_0 \;>\; |F(p)| - 2\varepsilon_0 \;>\; r - 2\varepsilon_0 \;>\; r - \varepsilon_3,$$

where initially $\varepsilon_0 < \varepsilon_3/2$.

(ii) Let $p \in D(p_j,\delta)\setminus(\cup_{k=1}^n Q_k(L_0))$. In this case we shall use (c1), (b1), (b6), (a2), (b7) and ε_0 is chosen less than $\varepsilon_3/5$

$$\begin{aligned}
|G(p)| &> |h_n(p)|-\varepsilon_0 \;>\; |h_j(p)|-2\varepsilon_0 \;>\; |h_{j-1}(p)+\kappa\,\zeta_j(p)e_1^j|-3\varepsilon_0\\
&\geq |{h_{j-1}}_{(1,S_j)}(p)+\kappa\,\zeta_j(p)|-3\varepsilon_0 \;>\; |F_{(1,S_j)}(p)+\kappa\,\zeta_j(p)|-4\varepsilon_0\\
&\geq |F_{(1,S_j)}(p_j)+\kappa\,\zeta_j(p)|-5\varepsilon_0 \;=\; ||F(p_j)|+\kappa\,\zeta_j(p)|-5\varepsilon_0\\
&> r-5\varepsilon_0 \;>\; r-\varepsilon_3.
\end{aligned}$$

(iii) Let $p \in int(D(p_j,\delta))\cap Q_j(L_0)$. First, we observe

$$\begin{aligned}
|h_n(a_j)-(F(p_j)+3(R-r)e_1^j)| &\leq |h_n(a_j)-h_j(a_j)|+|F(a_j)-F(p_j)|\\
&+ |h_j(a_j)-F(a_j)-3(R-r)e_1^j| \;<\; 3\varepsilon_0,
\end{aligned}$$

where we have used that $h_j(a_j)=h_{j-1}(a_j)+3(R-r)e_1^j$, and the formulas (a2) and (b1).

In addition, we have $|h_n(a_j)|>2R-r>R$ from (b4), and $|F(p_j)|+3(R-r)>3R-2r>R$.

Hence, as $F(p_j)+3(R-r)e_1^j=(|F(p_j)|+3(R-r))e_1^j$, one has

$$\begin{aligned}
\left|\overline{e}_2^j-e_1^j\right| &= \left|\frac{h_n(a_j)}{|h_n(a_j)|}-e_1^j\right| \;<\; 6\varepsilon_0 \min\left\{\frac{1}{|h_n(a_j)|},\frac{1}{|F(p_j)|+3(R-r)}\right\}\\
&< \frac{6\varepsilon_0}{R}.
\end{aligned}\tag{7}$$

Now, for the point p, we get

$$\begin{aligned}
|G(p)| &> |g_j(p)|-\varepsilon_0 \;\geq\; |{g_j}_{(2,\overline{S}_j)}(p)|-\varepsilon_0 \;>\; |{h_n}_{(2,\overline{S}_j)}(p)|-2\varepsilon_0\\
&> |{h_j}_{(2,\overline{S}_j)}(p)|-3\varepsilon_0 \;=\; |\langle h_j(p),\overline{e}_2^j\rangle|-3\varepsilon_0\\
&= |\langle h_j(p),e_1^j\rangle+\langle h_j(p),\overline{e}_2^j-e_1^j\rangle|-3\varepsilon_0\\
&\geq |{h_j}_{(1,S_j)}(p)|-|\langle h_j(p),\overline{e}_2^j-e_1^j\rangle|-3\varepsilon_0 \;>\; |{h_j}_{(1,S_j)}(p)|-\frac{6\varepsilon_0}{R}|h_j(p)|-3\varepsilon_0,
\end{aligned}$$

where we have used (c1) and (b1) again.

So, we estimate $|{h_j}_{(1,S_j)}(p)|$ and $|h_j(p)|$,

$$\begin{aligned}
|{h_j}_{(1,S_j)}(p)| &= |{h_{j-1}}_{(1,S_j)}(p)+\kappa\,\zeta_j(p)| \;>\; |F_{(1,S_j)}(p)+\kappa\,\zeta_j(p)|-\varepsilon_0\\
&> |F_{(1,S_j)}(p_j)+\kappa\,\zeta_j(p)|-2\varepsilon_0 \;=\; ||F_{(1,S_j)}(p_j)|+\kappa\,\zeta_j(p)|-2\varepsilon_0\\
&> |F(p_j)|-2\varepsilon_0 \;>\; r-2\varepsilon_0.
\end{aligned}$$

Here, we used (b1), (a2) and (b7).

Moreover,

$$|h_j(p)| \;\le\; |h_{j-1}(p)| + \kappa|\zeta_j(p)| + \varepsilon_0 \;<\; |F(p)| + 3(R-r) + 2\varepsilon_0 \;<\; 4R - 3r + 2\varepsilon_0,$$

from (b6), (b1) and (b7).

Therefore,

$$|G(p)| \;>\; (r - 2\varepsilon_0) - \frac{6\varepsilon_0}{R}(4R - 3r + 2\varepsilon_0) - 3\varepsilon_0 \;>\; r - \varepsilon_3.$$

Again, we have initially taken ε_0 small enough in terms of R, r and ε_3.

(iv) Let $p \in int(D(p_{j+1}, \delta)) \cap Q_j(L_0)$. This case is analogous to the previous one, bearing in mind

$$|\overline{e}_2^j - e_1^{j+1}| \;\le\; |\overline{e}_2^j - e_1^j| + |e_1^j - e_1^{j+1}| \;<\; \frac{6\varepsilon_0}{R} + \frac{\varepsilon_0}{3(R-r)}.$$

The last inequality is a consequence of (7) and (a3).

(v) Let $p \in Q_j(L_0) \setminus (\cup_{k=1}^n int(D(p_k, \delta)))$. From (c1), (b1), (7) and (a2), one obtains

$$\begin{aligned} |G(p)| \;&>\; |g_j(p)| - \varepsilon_0 \;\ge\; |g_{j(2,\overline{S}_j)}(p)| - \varepsilon_0 \;=\; |g_{j-1(2,\overline{S}_j)}(p)| - \varepsilon_0 \\ &>\; |h_{n(2,\overline{S}_j)}(p)| - 2\varepsilon_0 \;>\; |F_{(2,\overline{S}_j)}(p)| - 3\varepsilon_0 \;>\; |F_{(1,S_j)}(p)| - 3\varepsilon_0 - \frac{6\varepsilon_0}{R} \\ &>\; |F_{(1,S_j)}(p_j)| - 4\varepsilon_0 - \frac{6\varepsilon_0}{R} \;>\; r - 4\varepsilon_0 - \frac{6\varepsilon_0}{R} \;>\; r - \varepsilon_3. \end{aligned}$$

This finishes Lemma 3 and also proves Lemma 1. ☐

References

[AFM] A. Alarcón, L. Ferrer and F. Martín, Density theorems for complete minimal surfaces in $\mathbb{R}^3$, *Geom. Func. Anal.* **18** (2008), 1–49.

[CR] P. Collin and H. Rosenberg, Construction of harmonic diffeomorphisms and minimal graphs. *Preprint (arXiv:math.DG/0701547v1).*

[GR] J.A. Gálvez and H. Rosenberg, Minimal surfaces and harmonic diffeomorphisms from the complex plane onto a Hadamard surface. *Preprint (arXiv:math.DG/0807.0997v1).*

[G] A. Grigor'yan, Analytic and geometric background of recurrence and non-explosion of the brownian motion on Riemannian manifolds, *Bull. Am. Math. Soc* **36** (1999), 135–249.

[H] E. Heinz, Über die Lösungen der Minimalflächengleichung, *Nachr. Akad. Wiss. Göttingen. Math.-Phys. Kl.* (1952), 51–56.

[M] S. Morales, On the existence of a proper minimal surface in $\mathbb{R}^3$ with a conformal type of disk, *Geom. Funct. Anal.* **13** (2003), 1281–1301.

[R] H.L. Royden, Function theory on compact Riemann surfaces, *J. Anal. Math.* **18** (1967), 295–327.

[SY] R. Schoen and S.T. Yau, *Lectures on harmonic maps.* Conference Proceedings and Lecture Notes in Geometry and Topology. International Press, Cambridge MA, 1997.

[W] E.F. Wold, Embedding Riemann surfaces properly into $\mathbb{C}^2$, *Internat. J. Math.* **17** (2006), 963–974.

The first author is partially supported by MCYT-FEDER, Grant No MTM2007-61775 and the second author by MCYT-FEDER, Grant No MTM2007-65249. The authors are partially supported by Grupo de Excelencia P06-FQM-01642 Junta de Andalucía.